

Meier T., Hensen I. & Partzsch M. (2022) Effects of nitrogen addition and above-ground biomass removal on the growth and interactions between species of xerothermic grasses. – Preslia 94: 607–629.

Supplementary Table S1. – First-year and second year performance in terms of above-ground biomass of *Bromus erectus*, *Brachypodium pinnatum*, *Stipa capillata* and *S. tirsia* in the different nutrient conditions and clipping treatments.

	<i>B. erectus</i>	<i>B. pinnatum</i>	<i>S. capillata</i>	<i>S. tirsia</i>
first year				
N-	17.80 ± 3.47	12.79 ± 2.19	1.06 ± 0.38	4.43 ± 1.07
N+	13.97 ± 2.71	4.15 ± 1.54	0.93 ± 0.46	1.58 ± 0.33
second year				
N-, unclipped	20.27 ± 4.92	16.73 ± 2.63	0.24 ± 0.08	10.58 ± 1.74
N-, clipped	13.33 ± 2.11	9.56 ± 1.59	0.37 ± 0.11	6.40 ± 1.31
N+, unclipped	69.48 ± 9.85	7.55 ± 1.86	0.73 ± 0.18	1.39 ± 0.62
N+, clipped	14.39 ± 3.58	11.34 ± 1.97	0.91 ± 0.17	0.90 ± 0.32