

Phytogeographic and taxonomic remarks on eleven species of cyanophilic lichens from Central Europe

Fytogeografické a taxonomické poznámky k 11 druhům cyanofilních lišejníků střední Evropy

Helene Czeika¹, Gerhard Czeika¹, Anna Guttová², Edit Farkas³, László Lőkös⁴ & Josef Halda⁵

¹Ottakringerstrasse 78/11, A-1170 Wien, Austria; ²Institute of Botany, Slovak Academy of Sciences, Dúbravská cesta 14, SK-845 23 Bratislava, Slovakia, e-mail: anna.guttova@savba.sk; ³Institute of Ecology and Botany, Hungarian Academy of Sciences, Alkotmány 2–4, H-2163 Vácraátót, Hungary, e-mail: efarkas@botanika.hu; ⁴Hungarian Natural History Museum, Könyves Kálmán krt 40, H-1087 Budapest, Hungary, e-mail: lokos@bot.nhmus.hu; ⁵Okresní muzeum Orlických hor, Jiráskova 2, CZ-516 01 Rychnov nad Kněžnou, Czech Republic, e-mail: halda@dobruska.cz

Czeika H., Czeika G., Guttová A., Farkas E., Lőkös L. & Halda J. (2004): Phytogeographic and taxonomical remarks on eleven species of cyanophilic lichens from Central Europe. – Preslia, Praha, 76: 183–192.

Comments on 11 species of cyanophilic lichens are presented. A new combination *Peccania cernohorskyi* is proposed, commented on and typified. *Anema nodulosum*, *A. prodigulum*, *Lempholemma intricatum*, *Leptogium ferax*, *Porocyphus rehmicus* and *Zahlbrucknerella calcarea* are reported from Slovakia for the first time, *Leptogium biatorinum* and *L. magnussonii* from Hungary, and *Anema prodigulum*, *Heppia adglutinata*, *Leptogium biatorinum* and *Psorotichia taurica* from the Czech Republic. *Leptogium cretaceum* is lectotypified.

Key words: Central Europe, *Collemataceae*, *Heppiaceae*, *Lichinaceae*, lichens, phytogeography, taxonomy

Introduction

Cyanophilic lichens, especially crustose ones, are described in various works from different periods as “the most intricate, and perhaps the least understood” (Crombie 1875) or simply “problematic” (Schultz & Büdel 2002). The information presented for 11 species from three families (*Collemataceae*, *Heppiaceae*, *Lichinaceae*) are intended to increase the descriptive information on “the very tips of phylogenetic branches“, without which phylogenetic studies would make little progress (Landrum 2001).

Material and methods

The specimens were either collected in the field by the authors or selected from herbaria when working on the revision, as indicated in the section ‘Specimens examined’. Acronyms of the collections follow Holmgren et al. (1990). Labels are full-scale cited. Descriptions of the species are based on these specimens. Each species entry is supplemented by a reference to its publication and basionym. Data on the species *Leptogium biatorinum* s. l. includes, moreover, a synonym and its lectotypification. Entry for the species *Peccania cernohorskyi* includes revised material from the countries, from which it is reported.

Results

Anema nodulosum (Nyl.) Forss.

Nova Acta R. Soc. Scient. Upsal., ser. 3, 13: 93, 1885. – *Collema nodulosum* Nyl., Acta Soc. Linn. Bordeaux 21: 266, 1857.

Description: Thalli black, cushion-like, diameter 3–4 mm, composed of small, nodulose-, globose- or club-like, \pm pruinose lobes. Rhizohyphae tufted. Apothecia present, diameter up to 0.5 mm, \pm sunken into the lobes, disc red-brown, thalline exciple very thin. Spores ellipsoid, unicellular, 8 per ascus, $10\text{--}13 \times 6\text{--}9 \mu\text{m}$.

The genus *Anema* is insufficiently studied from a biological and nomenclatoric viewpoint. Worldwide, 10 cosmopolitan species have been registered (Kirk et al. 2001). Only a few are included in modern works (see Wirth 1995, Moreno & Egea 1992, Clauzade & Roux 1985). There is more information on *A. nodulosum* in Forssell (1885), Ozenda & Clauzade (1970) or Clauzade & Roux (1985). The species occurs in France (Ozenda & Clauzade 1970, Clauzade & Roux 1985) from where the lectotype originated (Henssen & Jørgensen 1990), Algeria, Morocco and Spain (Moreno & Egea 1992). It has not been reported from Slovakia (see Pišút et al. 1998, Lackovičová 2003).

Specimens examined: **Slovakia:** Strážovské vrchy Mts: ad rupes calcareas supra Súfov, ca 400 m, 3. VI. 1922, leg. J. Suza (PRM, ut *A. decipiens*, fertile, rev. H. Czeika & G. Czeika); Muránska planina plateau: ad arcem Muráň, c. 950 m, in rupibus calcareis Cigánka, 21. V. 1929, leg. J. Suza (PRM, ut *A. decipiens*, rev. H. Czeika & G. Czeika, cum *Collema* sp., *Placynthium garovaglii*, *P. subradiatum*, *Synalissa symphorea*).

Anema prodigulum (Nyl.) Henssen

Lichenologist 22: 139, 1990. – *Omphalaria prodigula* Nyl., Flora 62: 353, 1879.

Description: Thalli composed of groups of blue-black, occasionally pruinose, up to 1 mm wide lobes of the same height, attached to the substrate by rhizohyphae. In contrast to those of *A. nodulosum* the tiny thalli do not form cushions. Sterile lobes nodular, those bearing apothecia of \pm clavate shape. Apothecia present, diameter up to 0.7 mm, often prominent, mostly one per lobe. Disc red-brown, thalline exciple broad. Spores ellipsoidal, unicellular, 8 per ascus, $11\text{--}15 \times 7\text{--}9 \mu\text{m}$.

This species might be confused with those of the genus *Psorotichia* (Moreno & Egea 1992), but the difference is evident when the anatomy is studied. Moreover, the surface of the squamules differs in the two genera. According to Ozenda & Clauzade (1970), it is uneven, but not as rough, granular or verrucose as in *Anema*. The species is known from France (Ozenda & Clauzade 1970, Clauzade & Roux 1985), Morocco, Spain (Moreno & Egea 1992) and North America (Schultz 2002, see also detailed description). It is not reported from Slovakia and the Czech Republic (see Pišút et al. 1998, Vězda & Liška 1999, Lackovičová 2003).

Specimens examined: **Slovakia:** Malá Fatra Mts: Malý Rozsutec, dolomit, 1931, leg. Z. Černohorský (PRM, ut *A. decipiens*, rev. H. Czeika & G. Czeika); **Czech Republic:** Moravia occid.: Ivančice, in rupibus conglomeratis permicis in valle fluvii Rokytná prope pag. Budkovice, alt. 250 m s. m., 14. XII. 1956, leg. A. Vězda (W, Lich. Bohem. Exs. no. 34, ut *A. notarisii*, rev. H. Czeika & G. Czeika).

Heppia adglutinata (Kremp.) A. Massal.

Geneac. Lich. 8, 1854. – *Lecanora adglutinata* Kremp., Flora 24: 675, 1851.

Description: Olive-green thalli are squamulose to peltate, diameter up to 4 cm. Lower surface delimited by a well developed cortical layer. Apothecia present (diameter to 2 mm), immersed, exposing brown-red, urceolate to plain discs. Spores unicellular, ellipsoidal (20–23 × 7–8 μm). Hymenia stain red with iodine.

This species, reintroduced into the literature by Henssen (1994), is distinct from *H. lutosa* with which it is frequently synonymized (see Gyelnik 1935, Egea 1989, Hafellner & Türk 2001). It is reported from Europe and North America (Henssen 1994, Guttová & Palice 2002, Büdel et al. 2002). So far, it is not known from the Czech Republic (see Vězda & Liška 1999).

Specimen examined: **Czech Republic:** Vápence u Švarcentalu [= Černý Důl, Krkonoše Mts], 20. VII. 1924, leg. A. Hilitzer (PRM, ut *H. virescens*, rev. A. Guttová).

Lempholemma intricatum (Arnold) Zahlbr.

Cat. Lich. Univ. 3: 22, 1925. – *Omphalaria intricata* Arnold, Flora 52: 254, 1869.

Description: Brown-black to black thalli clustered in mats (diameter up to 1 cm). Lobes cylindrical, several mm long, tangled, surface gently longitudinally wrinkled, apices not swollen. Apothecia absent. The specimens are identical to the British material from Kirkcudbright, Black Craig (BM).

The status of this species is not clear. Certain forms resemble well developed morphs of *Lempholemma botryosum* (Coppins et al. 1992). In that case characters of the thallus, lobes and spores are useful. *L. intricatum* does not form umbilicate thalli like *L. botryosum* and the lobes are up to 5 mm long, whereas those of *L. botryosum* do not exceed 1 mm and the spores are larger (10–14 × 8–11 μm) than those of the latter species (7–9 × 5–7 μm) (Coppins et al. 1992, Massalongo 1856).

The species is reported from Great Britain (Coppins et al. 1992), Germany (Scholz 2000), Norway (Jørgensen 1988), Austria (Hafellner & Türk 2001) and Switzerland (Henssen 1969). It has not been reported from Slovakia (see Pišút et al. 1998, Lackovičová 2003).

Specimens examined: **Slovakia:** Slovenský raj Mts: reservatio Piecky, Veľký vodopád, ad ripam rivi, 48°56'20"N, 20°21'50"E, ad saxa calcarea, alt. 600 m s. m., 6. VIII. 1998, leg. J. Halda, Z. Palice & Š. Bayerová (herb. J. Halda, JPH/2782, rev. A. Guttová); Malá Fatra Mts: in monte Malý Rozsutec, loco subhumido, ad saxa dolomitica, alt. 1300 m, 9. V. 1973, leg. I. Pišút (BRA, rev. A. Guttová); Belianske Tatry: Südhang des Skalné vráta, Kalkfelswand, c. 1400 m s. m., 19. VIII. 1999, leg. H. Czeika & G. Czeika (herb. H. Czeika & G. Czeika).

Leptogium biatorinum s. l.

Leptogium biatorinum (Nyl.) Leight., Lich. Fl. Gr. Brit., ed. 3: 25, 1879. – *Collema biatorinum* Nyl., Acta Soc. Linn. Bordeaux 21: 268, 1857.

Leptogium cretaceum (Sm.) Nyl., Acta Soc. Linn. Bordeaux 21: 24, 1857. – *Lichen cretaceus* Sm., Engl. Bot. XI: 738, 1800. Type: E.B. 738, Dr. Noëhden from Eton, ex herb. Turner (BM – **lectotype, designated here**).

Description: Thalli brown-black, or when growing in the shade of a bluish-grey colour, spreading over rock surfaces as ± a continuous crust. Composed of granules (juvenile stages) or irregularly shaped squamules with sublobate, often ascending margins. Apothe-

cia frequent, sessile (diameter 0.20–0.75 mm), with a prominent proper exciple. The habitus is variable, which is why the collective name *L. biatorinum* s.l. is used here. The character of particular features varies within the delimitations of the taxa listed above (see Smith & Sowerby 1800, Leighton 1879, Ozenda & Clauzade 1970, Blum et al. 1975, Clauzade & Roux 1985, Jørgensen 1994, Guttová & Palice 2002, Wirth 1995 for further information).

The duplicate of Nylander's original material of *L. biatorinum* (Nyl.) Leight. (*Collema biatorinum*, ad Parisiens, ex herb. Leighton, BM) is terricolous. The thalli are composed of granules, which form tiny lobulate squamules. The sessile apothecia are globular. The lower part of the thin thalline exciple proliferates locally forming squamule-like outgrowths comparable to those of e.g. *L. lichenoides*. If the original material (possibly in H. P. M. Jørgensen, pers. com.) is not located, this duplicate specimen could become the lectotype.

The original material of *Leptogium cretaceum* (Sm.) Nyl. consists of two saxicolous specimens (*Lichen cretaceus*): 1. Sowerby's herbarium Feb. 1859, 2. E.B. 738, Dr. Noëhden from Eton, ex herb. Turner (both BM). The former was studied by Henssen in 1976 and labelled as the type. It corresponds with illustrations published in the 11th volume of English Botany (Smith & Sowerby 1800). The thalli grow on flint stones and are composed of stellately radiating lobate squamules with sessile apothecia centrally. The latter is not so well developed and directly refers to the one described by Smith in the sentence "Our perfect specimens were gathered on chalk by Dr. Noehden, and Mr. Gotbed of Eton" (Smith & Sowerby 1800). We designate it as a lectotype.

This taxon is reported from many European countries and the USA (Hafellner & Türk 2001, Alstrup & Sjøchting 1989, Ozenda & Clauzade 1970, Scholz 2000, Coppins & Purvis 1992, Nimis 1993, Motiejūnaite et al. 1998, Motiejūnaite 1999, Diederich & Sérusiaux 2000, Faūtynowicz 1993, Santesson 1993, Kondratyuk et al. 1998, Jørgensen & Tønberg 1999). It has not been reported from the Czech Republic and Hungary (see Vězda & Liška 1999, Verseghe 1994).

Specimens examined: **Czech Republic:** Moravia, Brno: ad marginem silvae pr. p. Mokrá Hora, c. 300 m s. m., ad terram loessaceam, VI. 1960, leg. A. Vězda (W, ut *L. byssinum*, rev. H. Czeika & G. Czeika); Brno: inter pagos Soběšice et Mokrá Hora (prope urbem Brno), ad terram arenaceam et supra muscos in fossis viae sylvaticae, alt. 300 m s. m. 28. II. 1961, leg. A. Vězda (Lich. Sel. Exs. no 81, ut *L. subtile*) [BP, BRA; rev. A. Guttová], ibid. alt. 250 m, IV. 1966, leg. A. Vězda (BM, ut *L. byssinum*, rev. P. M. Jørgensen & A. Guttová); Bohemia septentr.: ad muros arenaceos humidos prope Příhrazy, Mladá Boleslav, VII. 1965, leg. Vězda (BRA, ut *L. cf. tenuissimum*, rev. A. Guttová); **Hungary:** Pilis Mts, "Visegrádi-hegység": Visegrád, the valley of the stream Apátkúti-patak (Apátkúti-völgy), mixed deciduous forest (*Carpinus betulus*, *Fagus sylvatica*, *Quercus* sp. div.) c. 2.5 km from Visegrád past abandoned quarry, andesite outcrops along the road bank, on shaded mossy andesite rock, alt. c. 350 m a.s.l., 21. XI. 2002, leg. A. Guttová, E. Farkas & L. Lőkös, fertile (SAV, BP).

Leptogium ferax (Dur. et Mont.) Rabenh.

Sitzungs-Ber. Naturwiss. Ges. Isis, Dresden, 1870: 237, 1871. – *Collema ferax* Dur. et Mont., Flore d'Algérie, Crypt. 1: 202, 1846.

Description: Thalli mostly brown, if shaded then of a greenish tinge, not exceeding 1 cm in diameter. Lobes orbicular, margin entire, lamina subtly wrinkled (as in e.g. *L. magnussonii*). Medulla compact, comparable to that of *L. magnussonii* (cf. Jørgensen 1994). Apothecia present, diameter 2–2.5 mm, rim lined with dense accumulation of scale- or shell-like structures recalling the isidia of *Collema flaccidum*. Disc plane red-

brown. Spores ellipsoidal (immature ones fusiform to broadly fusiform), submuriform to muriform, 8 per ascus, $30.9 (-32.2) \times 9.2 (-11.5) \mu\text{m}$.

This species resembles the common gelatinous lichen *Collema crispum* (Degelius 1954). A well developed cortex and muriform spores are diagnostic characters. It has been recorded from the coastal regions of Mediterranean Algeria, France, Greece (Crete), Portugal and Spain (Arvidsson 1984, Carvalho & Jones 1997). Despite its conspicuous habit the species is not reported from inland localities. The western slopes of the Malé Karpaty Mts (SW Slovakia) experience special climatic conditions, which is reflected in the presence of several submediterranean and pannonian vascular plants. Taking this phytogeographical aspect into account, we expected to find this species there but finding a new macrolichen in an area frequently visited by lichenologists (Lackovičová 1978, 1988) is certainly interesting. Nevertheless, the species has not been reported from Slovakia up to now (see Pišút et al. 1998, Lackovičová 2003).

Specimens examined: **Slovakia:** Malé Karpaty Mts: Plavecký Mikuláš, unnamed hill with limestone/conglomerate outcrops and short xerotherm vegetation above central part of the village (E slope of the elevation 367.2), on open, exposed calcareous soil in rock fissures among mosses (*Bryum argenteum*, *Tortella tortuosa*, *Hypnum cupressiforme*, *Tortula muralis*, *Encalypta vulgaris*, *Ditrichum flexicaule*), alt. c. 280 m a.s.l., 5. VI. 2001, leg. A. Guttová (7469d), fertile (SAV, together with *Agonimia opuntiella*, *Caloplaca* sp., *Mycobilimbia* sp.); Plavecký Mikuláš: Kršlenica national nature reserve, S exposed limestone/conglomerate ridge of cliffs, on soil among mosses (*Encalypta* cf. *vulgaris*), alt. c. 480 m a.s.l., 5. IV. 2002, leg. A. Guttová, A. Lackovičová, E. Lisická, J. Liška & V. Orthová (7469d), fertile (SAV); Plavecké Podhradie: Pohanská national nature reserve, W slope of the hill Pohanská, abandoned quarry currently used as shooting-range, on sandy soil among mosses (*Ceratodon purpureus*, *Didymodon* sp., *Tortula muralis*, *Encalypta streptocarpa*, *Fissidens* sp.), alt. c. 300 m a.s.l., 8. XI. 2002, leg. A. Guttová, V. Kučera, A. Lackovičová & V. Orthová (7569b), fertile (SAV, conf. P. M. Jørgensen XII. 2002).

Leptogium magnussonii Degel. et P. M. Jørg.

Lichenologist 26 (1): 14, 1994.

Description: Parts of the thalli brown to brown-black, 2–3 cm in diameter. Lobes rounded, lamina sparingly covered with granular or cylindrical isidia. Medulla compact. Apothecia absent.

Since 1994, when this isidiate *Leptogium* species was described (Jørgensen 1994), more information has been recorded on its distribution and ecological requirements (Berger 1996, Sérusiaux et al. 1999, Guttová 2000). This species is on the redlist in Sweden (Thor & Arvidsson 1999). The find recorded here is the first for Hungary (see Verseghy 1994, 2003).

Specimens examined: **Hungary:** Pilis Mts, Visegrádi-hegység: Kő-hegy Mt., on the rocks at Petőfipihenő N of Pomáz, on siliceous rock, alt. 350 m a.s.l., 19. V. 1985, leg. E. Farkas & L. Lőkös (VBI ut *L. lichenoides*, rev. P. M. Jørgensen ut *L. magnussonii* 29. I. 2001); Pilis Mts, Visegrádi-hegység: along the stream Apátkúti-patak in the valley Apátkúti-völgy) and on the hill Apátkúti-bérc 1–2 km S of Visegrád, on mossy soil, on siliceous rocks, alt. 350 m, 3. IV. 1988, leg. E. Farkas & L. Lőkös (VBI ut *L. lichenoides*, rev. P. M. Jørgensen ut *L. cf. magnussonii* 29. I. 2001).

Peccania cernohorskyi (Servít) Czeika et Guttová, **comb. nova**

Thyrea cernohorskyi Servít, Věstn. Král. České Společn. Nauk, ser. math.-nat., 1934: 6, 1935. Type: diabasy u sv. Ivana 25. IX. 1930 [Z. Černohorský & A. Hilitzer] (PRM – **lectotype, designated here**).

In 1930, Z. Černohorský and A. Hilitzer collected a diminutive terricolous/saxicolous cyanolichen at a locality in the Bohemian Karst [= Český kras] S of Prague, frequently visited by botanists. After examining the material Servít recognized it as an undescribed species of the genus *Thyrea* and named it after one of the collectors – “*Thyrea Černohorskýi*” Servít (see Servít & Černohorský 1935). His description is thorough, consisting of a Latin diagnosis and comprehensive German commentary on details of apothecial anatomy and a comparison with the species *Thyrea camaromorpha*. The text is supplemented by accurate illustrations of a fertile thallus (also in cross-section), asci with spores and paraphyses.

Almost 60 years later, the taxon “*Černohorskýi*” was revised by Henssen & Jørgensen (1990). They transferred *Thyrea cernohorskyi* Servít to the genus *Anema* [*A. cernohorskyi* (Servít) Henssen], although no original material was located.

In 2000, when revising different taxa of cyanolichens from Slovakia and the Czech Republic deposited in PRM, we located the original material and duplicates of *Thyrea cernohorskyi* in BRA and PRC. The material corresponds with the original description and illustrations (thallus surface not smooth, covered with spherical/verrucose outgrowths, apothecia diameter 1–1.5 mm, upper part of hymenium orange-brown, 8 spores/ascus, $12\text{--}15 \times 12\text{--}14 \mu\text{m}$). The thalli are heteromerous (growing either on a layer of soil/dust with bryophytes or directly on rock). They also bear pycnidia producing filiform conidia, (16) $19\text{--}25 \times \pm 1 \mu\text{m}$. The hymenium is delimited exclusively by a thalline exciple. Recent material collected by J. Kocourková at the same locality (PRM) is identical with the original material, but sterile, producing only filiform conidia.

In terms of the recent delimitation of the genus *Anema* (Henssen & Jørgensen 1990, Moreno & Egea 1992, Schultz & Büdel 2002) the following characters are crucial: thallus squamulose/subfruticulose/peltate, homoiomerous, lacking central hyphal strand, fruiting bodies pycnoascocarps, hymenium colourless, thalline exciple well developed, proper exciple inconspicuous or missing, conidia ellipsoid. Not all characters of the thalli, however, correspond to this description. The habit is subfruticulose to peltate (see Servít & Černohorský 1935: “ex initiis globosis monophyllinus, demum p.p. crustosus... thalli monophylli 2–3 mm lati, irregulariter subcrenulati vel lobulati, subtus late umbilicati...”) and thalli are not fixed to the substrate by a central holdfast, but attached at several spots. The conidia are filiform as in the genus *Peccania*, representatives of which are the only ones with filiform conidia in the related groups *Anema* – *Thyrea* – *Peccania* (Moreno & Egea 1992). For this reason, the combination *Peccania cernohorskyi* (Servít) Czeika et Guttová is proposed. As there are several specimens collected at the same time in different herbaria, a lectotype is designated. The specimens in BRA and PRC are isolectotypes.

Specimens examined: **Czech Republic:** Diabasy u sv. Ivana [= Svatý Jan pod Skalou], ad saxa diabasica prope sv. Ivan, 25. IX. 1930 [Z. Černohorský & A. Hilitzer] (PRM, ut *Thyrea Černohorskýi*; BRA, rev. A. Guttová, H. Czeika & G. Czeika), ibid. saxa diabasica prope sv. Ivan, 25. IX. 1930, A. Hilitzer (BRA, ut *Thyrea Černohorskýi*, rev. A. Guttová); ibid., 25. IX. 1930, leg. Z. Černohorský (PRC, ut *Thyrea Černohorskýi*, rev. A. Guttová); ibid., 25. IX. 1998, leg. J. Kocourková (PRM, rev. A. Guttová, H. & G. Czeika).

Besides the Czech Republic, *Anema cernohorskyi* (Servít) Henssen is recorded from Germany (Scholz 2000). The entry is based upon Henssen: Lich. Cyan. Fungi Sax. Exs. no 47 (BRA, W; rev. A. Guttová, H. Czeika & G. Czeika). However, the material in this collection is *Anema tumidulum* Henssen ined., like that recorded by Santesson (1993) from Norway, from calcareous rock in Buskerud and Telemark.

Specimens examined: **Norway:** Buskerud, Hole, the island Storøya in lake Tyrifjorden, south-western part, alt. c. 65 m, calcareous rock face, 14. XI. 1981, leg. E. Timdal no 3142 (O, rev. A. Guttová); Buskerud, Hole, Purkøya, the SW-side, alt. 63–75 m, on calcareous rock, steep rock wall, 8. IX. 1997, leg. H. Bratli & E. Timdal no 8720 (O, rev. A. Guttová); Buskerud, Hole, Burudåsen, between Kongelv and Kleiva, alt. 90 m, steep, W-facing, calcareous rock wall in edge of pine forest, 29. I. 1989, leg. J. Holtan-Hartwig & E. Timdal (O, rev. A. Guttová); Buskerud, Hole, limestone cliffs c. 500 m SE of Bønsnes church, alt. 65–80 m, 9. IV. 1994, leg. R. Haugan & E. Timdal no 7873 (O, rev. A. Guttová); Buskerud, Hole, cliffs S of Stammes at the shore of lake Tyrifjorden, alt. 70 m, steep, SE-facing cliffs of calcareous rock, 29. I. 1989, leg. J. Holtan-Hartwig & E. Timdal (O, rev. A. Guttová); Buskerud, Hole, Lemostangen in lake Tyrifjorden, alt. 65 m, 31. X. 1981, leg. E. Timdal no 3105 (O, rev. A. Guttová); Telemark, Bamble, Langesundstangen, the west side, alt. 1–40 m, on vertical, calcareous rock wall, 22. VI. 1997, leg. H. Bratli & E. Timdal nos 1096, 8738 (O, rev. A. Guttová); Telemark, Porsgrunn, Kotøya, alt. 2–5 m, steep to vertical rock wall, facing west, calcareous rock, 8. VI. 1996, leg. H. Bratli & E. Timdal (O, rev. A. Guttová).

Porocyphus rehmicus (A. Massal.) Zahlbr.

Cat. Lich. Univ. 2: 756, 1924. – *Psorotichia rehmica* A. Massal., *Miscell. Lichenol.* 53, 1865.

Description: Thallus black, granular, non-areolate, photobiont is a filamentous cyanobacterium (*Calothrix* sp.). Apothecia present, discs remain pore-like. Paraphyses shortly branched, apices not swollen. Spores both ellipsoidal (12.5–15 × 5–7.5 µm) and spherical.

The thallus corresponds to that described for *P. rehmicus*, although this species is reported to prefer calcareous substrates (Migula 1929, Henssen 1963, Ellis 1981, Moreno & Egea 1994). According to others the thalli bearing this name could be an aberrant form of *P. coccodes*, growing on silicate (Coppins & Gilbert 1992). Despite all these uncertainties, the crucial character distinguishing the two species is the shape of the thallus; in *P. rehmicus* it is granular or minutely shrubby and in *P. coccodes* areolate.

The species might have been overlooked in Slovakia from where it has not been reported (see Pišút et al. 1998, Lackovičová 2003).

Specimen examined: **Slovakia:** Malá Fatra Mts: auf einem Granitfelsen zwischen Kralován und Párnica, Com. Árva in Ungarn, 17. VII. 1883, leg. Lojka (Gránitsziklán Kralován és Párnica Között. Árvam.), ut *Psorotichia* (BP, rev. A. Guttová).

Psorotichia taurica (Nyl.) Vain.

Termész. Füzet. 22: 312, 1899. – *Collemopsis taurica* Nyl. *Flora* 64, p. 97, 1886.

Description: Thallus black, brown-black, forming irregular patches of diameter 1–3 cm, growing preferably in rock fissures and caverns. Composed of granules to irregularly shaped areoles (diameter up to 0.5 mm), thin (up to 0.3 mm). Apothecia ± abundant, diameter up to 0.25 mm, globose. Disc concolorous with thallus, urceolate, exposed mostly through a pore-like opening. Epithemium and upper part of hymenium emerald green or blue-green. Spores predominantly spherical (diameter 7–8 µm), exceptionally broadly ellipsoidal (7–10 × 6–8 µm), 16 per ascus. For additional information consult Nylander (1886) and Oksner (1956).

The type of the species is reported from the Crimean peninsula, Ukraine (Nylander 1886, Oksner 1956, Kondratyuk et al. 1998). Kopatchevskaya (1986) regarded this species as an element of the European area, Crimean-Caucasian group. It has not been found in the Czech Republic (see Vězda & Liška 1999).

Specimens examined: **Czech Republic:** Central Bohemia, Praha–Hlubočepy: the Prokopské údolí valley, an abandoned flooded quarry, on steep, periodically inundated Ca-rock above the small lake, alt. 240 m a.s.l., 18. IV. 1999, leg. Z. Palice, L. Bičanová & K. Vincencová (herb. Z. Palice no. 1879, det. H. Czeika); *ibid.* 25. VI. 2002, leg. A. Guttová & Z. Palice (SAV).

Zahlbrucknerella calcarea (Herre) Herre

J. Wash. Acad. Sc. 2: 384, 1912. – *Zahlbrucknera calcarea* Herre, Proc. Wash. Acad. Sci. 12: 129, 1910.

Description: The black thalli are up 1 cm in diameter, composed of tufts of fine filaments, which nearly completely cover the photobiont (*Scytonema* sp.) and the mycobiont is concentrated mostly in conspicuous hyaline basal strand. Apothecia absent.

When fertile, the species is easy to identify because of the peculiar polysporous asci (16–24 spores – Henssen 1977, Kantvilas & Elix 1992). Despite it having a cosmopolitan distribution, it is rarely recorded. This, to a certain extent, might be due to its ecological requirements. It prefers the bases of perpendicular rock-faces, where collecting is often difficult. It has not been reported from Slovakia (see Pišút et al. 1998, Lackvičová 2003).

Specimens examined: **Slovakia:** Strážovské vrchy Mts: ad rupes calcareas supra Súľov, c. 500 m, 3. VI. 1922, leg. J. Suza (PRM, ut *A. decipiens*, rev. H. & G. Czeika, cum *Synalissa symphorea*, *Anema tumidulum*, *Placynthium subradiatum*, *Collema* sp.); Tatry Mts (Bélai mészalpok) – mészkősziklafalon a Greiner-en, c. 2000 m, Szepes vm, 9. VIII. 1916, leg. Timkó (BP, cum *Placynthium coeruleascens*); Belianske Tatry Mts: Südhang des Skalné vráta, Kalkfelswand, c. 1400 m s. m., 19. VIII. 1999, leg. H. Czeika & G. Czeika (herb. H. Czeika & G. Czeika).

Acknowledgement

We thank P. M. Jørgensen (Bergen), G. Kantvilas (Hobart), K. Marhold (Bratislava), Z. Palice (Praha) and I. Pišút (Bratislava) for valuable comments, J. Kocourková (Praha) who kindly provided us with her collections of cyanolichens, K. Kresáňová (Bratislava) for determination of bryophytes and the curators of the cited herbaria for their assistance. Tony Dixon (Norwich) kindly improved our English. The work was supported by grant projects VEGA 4035/04, APVT 51–005102 and OTKA T 047/60.

Souhrn

Príspevek obsahuje komentáre k fytoogeografii, prípadne k taxonomii 11 druhů cyanofilních lišejníků z čeledí *Collemataceae*, *Heppiaceae* a *Lichinaceae*. Je navržena nová kombinace *Peccania cernohorskyi*, doplněná typifikací. Druhy *Anema nodulosum*, *A. prodigulum*, *Lempholemma intricatum*, *Leptogium ferax*, *Porocyphus rehmicus* a *Zahlbrucknerella calcarea* jsou udávány poprvé z území Slovenska, *Leptogium biatorinum* a *L. magnussonii* poprvé z Maďarska. Z území České republiky jsou poprvé udávány následující taxony: *Anema prodigulum*, *Heppia adglutinata*, *Leptogium biatorinum* a *Psorotichia taurica*. Pro druh *Leptogium cretaceum* byl vybrán a ustanoven lektotyp.

References

- Alstrup V. & Søchting U. (1989): Checkliste og status over Danmarks laver. – Nordisk Lichenologisk Forening, København. [44 pp.]
- Arvidsson L. (1984): Two new records of *Leptogium ferax*. – Lichenologist 16: 91–92.
- Berger F. (1996): Neue und seltene Flechten und lichenicole Pilze aus Oberösterreich, Österreich II. – Herzogia 12: 45–84.
- Blum O. B., Dombrovskaya A. V., Inashvili C. N., Piterans A. V., Roms E. G. & Savicz V. P. (1975): Handbook of the lichens of the U.S.S.R. 3. *Caliciaceae* – *Gyalectaceae*. – Nauka, Leningrad. [275 pp.]
- Büdel B., Schultz M. & Nash T. H. III (2002): *Heppia*. – In: Nash T. H. III, Ryan B. D., Gries C. & Bungartz F. (eds), Lichen flora of the Greater Sonoran Desert region. Vol. 1 (the pyrenolichens and most of the squamulose and macrolichens), p. 97–98, Lichens Unlimited, Arizona State Univ., Tempe.

- Carvalho P. & Jones M. P. (1997): New and interesting lichens from Portugal. – *Cryptogamie, Bryologie, Lichénologie* 18: 291–294.
- Clauzade G. & Roux C. (1985): Likenoj de okcidenta Europo, ilustrita determinlibro. – *Bull. Soc. Bot. Centre-Ouest, sér. n., num. spéc.* 7: 1–893.
- Coppins B. J. & Gilbert O. L. (1992): *Porocyphus*. – In: Purvis O. W., Coppins J. B., Hawksworth D. L., James P. W. & Moore D. M. (eds.), *The lichen flora of Great Britain and Ireland*, p. 492–493, *Nat. Hist. Mus. Publ. & British Lichen Soc.*, London.
- Coppins B. J., Gilbert O. L. & Jørgensen P. M. (1992): *Lempholemma*. – In: Purvis O. W., Coppins B. J., Hawksworth D. L., James P. W. & Moore D. M. (1992): *The lichen flora of Great Britain and Ireland*, p. 341–344, *Nat. Hist. Mus. Publ. & British Lichen Soc.*, London.
- Coppins B. J. & Purvis O. W. (1992): *Leptogium*. – In: Purvis O. W., Coppins B. J., Hawksworth D. L., James P. W. & Moore D. M. (eds.), *The lichen flora of Great Britain and Ireland*, p. 350–356, *Nat. Hist. Mus. Publ. & British Lichen Soc.*, London.
- Crombie J. M. (1875): Revision of the British Collemacei. – *Grevillea* 3: 92–95.
- Degelius G. (1954): The lichen genus *Collema* in Europe. – *Symb. Bot. Upsal.* 13/2: 1–499.
- Diederich E. & Sérusiaux E. (2000): The lichens of Belgium and Luxembourg. An annotated checklist. – *Muséé National d'Histoire Naturelle, Luxembourg*. [207 pp.]
- Egea J. M. (1989): Los géneros *Heppia* y *Peltula* (líquenes) en Europa Occidental y Norte de Africa. – *Bibl. Lichenol.* 31: 1–122.
- Ellis L. T. (1981): A revision and review of *Lemmopsis* and some related species. – *Lichenologist* 13: 123–139.
- Fałtynowicz W. (1993): A checklist of Polish lichen forming and lichenicolous fungi including parasitic and saprophytic fungi occurring on lichens. – *Polish Bot. Stud.* 6: 1–65.
- Forsell K. B. J. (1885): Anatomie und Systematik der Gloeolichenen. – *Nova Acta Reg. Soc. Sci. Upsaliensis, ser. 3*, 13: 1–118.
- Guttová A. (2000): Three *Leptogium* species new to Central Europe. – *Lichenologist* 32: 291–303.
- Guttová A. & Palice Z. (2002): Lišajníky Národného parku Muránska planina II – Javorníková dolina. – *Výskum a ochrana prírody Muránskej planiny* 3: 53–68.
- Gyelnik V. (1935): De familiam Heppiacearum. I. – *Feddes Repert.* 38: 153–157.
- Hafellner J. & Türk R. (2001): Die lichenisierten Pilze Österreichs – eine Checkliste der bisher nachgewiesenen Arten mit Verbreitungsangaben. – *Stappia* 76: 1–167.
- Henssen A. (1963): Eine Revision der Flechtenfamilien *Lichinaceae* und *Ephebeaceae*. – *Symb. Bot. Upsal.* 18/1: 1–123.
- Henssen A. (1969): *Lichenes cyanophili exsiccati. Fasc. I (Nos. 1–25)*. – *Bot. Inst. Univ. Marburg*. [6 pp.]
- Henssen A. (1977): The genus *Zahlbrucknerella*. – *Lichenologist* 9: 17–46.
- Henssen A. (1994): Contribution to the morphology and species delimitation in *Heppia* sensu stricto (lichenized Ascomycotina). – *Acta Bot. Fenn.* 150: 57–73.
- Henssen A. & Jørgensen P. M. (1990): New combinations and synonyms in the *Lichinaceae*. – *Lichenologist* 22: 137–147.
- Holmgren P. K., Holmgren N. H. & Barnett L. (1990): *Index herbariorum. Part I: The herbaria of the world*. – *Regnum Veg.* 120: 1–693.
- Jørgensen P. M. (1988): Nyfunn av lav med blågrønne bakterier i Norge. – *Graphis Scripta* 2: 55–59.
- Jørgensen P. M. (1994): Further notes on European taxa of the lichen genus *Leptogium*, with emphasis on the small species. – *Lichenologist* 26: 1–29.
- Jørgensen P. M. & Tønsberg T. (1999): Notes on some small species of *Leptogium* from Pacific North America. – *The Bryologist* 102: 412–417.
- Kantvilas G. & Elix J. A. (1992): A new species and new records from the Tasmanian lichen flora. – *Muelleria* 7: 507–517.
- Kirk P. M., Cannon P. F., Davis J. C. & Stalpers J. A. (2001): *Ainsworth & Bisby's dictionary of the fungi*. – C.A.B. International, Wallingford [624 pp.]
- Kondratyuk S. Ya., Khodosovtsev A. Ye. & Zelenko S. D. (1998): The second checklist of lichen forming, lichenicolous and allied fungi of Ukraine. – *Phytosociocentre, Kiev*. [179 pp.]
- Kopatchevskaya E. G. (1986): Lichenoflora Kryma i jejo analiz. – *Naukovaya dumka, Kiev*. [294 pp.]
- Lackovičová A. (1978): Lišajníky Malých Karpát. – *Acta Ecol.* 6 (1977): 7–107.
- Lackovičová A. (ed.) (1988): *Dr. Alexander Zahlbruckner (1860–1938). Osobnosť a dielo*. – *Múzeum Petra Jilemnického, Jur pri Bratislave, Bratislava*. [60 pp.]
- Lackovičová A. (2003): Súhrn doplnkov k zoznamu a bibliografii lišajníkov Slovenska. – *Bull. Slov. Bot. Spoloč.* 25: 17–29.

- Landrum L. R. (2001): What has happened to descriptive systematics? What would make it thrive? – *Syst. Bot.* 26: 438–442.
- Leighton W. A. (1879): The lichen flora of Great Britain, Ireland, and the Channel Islands. Ed. 3. – Shrewsbury. [547 pp.]
- Massalongo A. (1856): *Miscellanea lichenologica*. – Tip. G. Civelli, Verona & Milano, p. 31–87.
- Migula W. (1929): Kryptogamen-Flora von Deutschland, Deutsch-Österreich und der Schweiz im Anschluss an Thomés Flora von Deutschland. Band 4. Flechten (1. Teil). Aufbau und Fortpflanzung des Flechtenkörper-Biologie der Flechten-Gymnocarpeae Scheibenfrüchtige. Unterreihe *Cyclocarpineae*, Familie *Physciaceae* bis *Gyrophoraceae*. – In: Thomé O. W., Flora von Deutschland, Österreich und der Schweiz in Wort und Bild, Band 11, Abt. 1, p. 1–527, H. Bermühler Verlag, Berlin-Lichterfelde.
- Moreno P. P. & Egea J. M. (1992): Estudios sobre el complejo *Anema-Thyrea-Peccania* en el sureste de la península Iberica y norte de Africa. – *Acta Bot. Barcinonensia* 41: 1–66.
- Moreno P. P. & Egea J. M. (1994): El género *Psorotichia* y especies próximas en el sureste de España y norte de Africa. – *Bull. Soc. Linn. Provence* 45: 291–308.
- Motiejūnaite J. (1999): Checklist of lichens and allied fungi of Lithuania. – *Bot. Lithuan.* 5: 251–269.
- Motiejūnaite J., Nordin A., Zalewska A., Bjelland T., Hedenas H., Westberg M., Heidmarsson S. & Prigodina I. (1998): Materials on lichen and allied fungi of Neringa National Park (Lithuania). – *Bot. Lithuan.* 4: 285–305.
- Nimis P. L. (1993): The lichens of Italy. – Museo Regionale di Scienze Naturali, Torino. [987 pp.]
- Nylander W. (1886): Addenda nova ad Lichenographiam europaeam. – *Flora* 64: 97–101.
- Oksner A. M. (1956): Flora lishainikiv Ukraini. Tom 1. – Vidav. Akad. Nauk Ukrainskoi RCR, Kiev. [495 pp.]
- Ozenda P. & Clauzade G. (1970): Les Lichenes. – Masson & Cie, Paris. [801 pp.]
- Pišút I., Guttová A., Lackovičová A. & Lisická E. (1998): Lichenizované huby (lišajníky). – In: Marhold K. & Hindák F. (eds.), Zoznam nižších a vyšších rastlín Slovenska, p. 229–295, Veda, Bratislava.
- Santesson R. (1993): The lichens and lichenicolous fungi of Sweden and Norway. – SBT-förlaget, Lund. [240 pp.]
- Scholz P. (2000): Katalog der Flechten und flechtenbewohnenden Pilze Deutschlands. – *Schr. Vegetationskd.* 31: 1–298.
- Schultz M. (2002): *Anema*. – In: Nash T. H. III, Ryan B. D., Gries C. & Bungartz F. (eds), Lichen flora of the Greater Sonoran Desert region. Vol. 1 (the pyrenolichens and most of the squamulose and macrolichens), p. 47, Lichens Unlimited, Arizona State Univ., Tempe.
- Schultz M. & Büdel B. (2002): Key to the genera of the *Lichinaceae*. – *Lichenologist* 34: 39–62.
- Sérusiaux E., Diederich P., Brand A. M. & van den Boom P. (1999): New or interesting lichens and lichenicolous fungi from Belgium and Luxembourg. VIII. – *Lejeunia* 162: 1–95.
- Servít M. & Černohorský Z. (1935): Flechten aus Čechoslovakiei. IV. Malá Fatra, Veľká Fatra und Choč-Gruppe in Slowakei nebst Nachträgen. – *Věstn. Král. České Společn. Nauk, ser. math.-nat.*, 1931: 1–34.
- Smith J. E. & Sowerby J. (1800): *English botany*. Vol. 11. – Davis, Wilks & Taylor, London.
- Thor G. & Arvidsson L. (eds.) (1999): Rödlistade lavar i Sverige – Artfakta. – ArtDatabanken, SLU, Uppsala. [528 pp.]
- Verseghy K. (1994): Magyarország zuzmóflórájának kézikönyve. – *Studia Naturalia* 5, Magyar Természettudományi Múzeum, Budapest. [415 pp.]
- Vězda A. & Liška J. (1999): Katalog lišejníků České republiky. – Botanický ústav AV ČR, Průhonice. [283 pp.]
- Wirth V. (1995): Die Flechten Baden-Württembergs. Teil 1, 2. – Ulmer, Stuttgart. [1006 pp.]

Received 25 September 2003

Revision received 16 February 2004

Accepted 7 April 2004